arXiv:1211.5369v1 [math.PR] 22 Nov 2012

Cauchy-Laguerre two-matrix model and the Meijer-G random point field

M. Bertola†, M. Gekhtman‡, J. Szmigielski ⋆

† *CRM, Université de Montréal and Department of Mathematics and Statistics, Concordia University*

‡ *Department of Mathematics, University of Notre Dame*

⋆ *Department of Mathematics and Statistics, University of Saskatchewan*

Abstract

We apply the general theory of Cauchy biorthogonal polynomials developed in [2],[3] to the case associated with Laguerre measures. In particular, we obtain explicit formulæ in terms of Meijer-G functions for all key objects relevant to the study of the corresponding biorthogonal polynomials and the Cauchy two-matrix model associated with them. The central theorem proved for such Cauchy two-matrix model is that a scaling limit of the correlation functions for eigenvalues near the origin exists and is given by certain new random point field, *the Meijer-G random field*. We conjecture that this random point field leads to a novel universality class of random fields parametrized by exponents of Laguerre weights.

Contents

1 Introduction 2
- 1.1 Summary of results . . . 3
  - 1.1.1 The scaling random field at the origin: the Meijer-G random point field . . . 4
  - 1.1.2 Outlook . . . 5

2 Setup 6
- 2.1 The kernels of the correlation functions . . . 7
- 2.2 Facts on Jacobi Polynomials . . . 8

3 Cauchy-Laguerre Biorthogonal polynomials 11
- 3.1 The auxiliary functions $p_n^{(1)}$ and $p_n^{(2)}$ . . . 12
- 3.2 The kernels for finite n . . . 15
- 3.3 The Meijer-G random point field . . . 17
  - 3.3.1 Asymptotics of the kernels $K_{\mu\nu}^{(n)}$ . . . 18
  - 3.3.2 Expressions of the kernels in “integrable” form . . . 20

A The Meijer-G functions 22

B Correlation functions **23**

1 Introduction

The Cauchy two-matrix model was introduced in [2] as a new kind of an integrable multi matrix model. The model consists of a probability measure on the space of pairs M_1, M_2 of $n \times n$ *positive semidefinite* Hermitean matrices. This probability measure depends on the choice of two scalar functions $V_1, V_2 : \mathbb{R}_+ \to \mathbb{R}$, called the *potentials* and is defined as

$$\mathrm{d}\mu(M_1, M_2) = \frac{1}{\mathcal{Z}_n} \frac{\mathrm{e}^{-N\mathrm{Tr}(V_1(M_1)+V_2(M_2))}}{\det(M_1+M_2)^n} \mathrm{d}M_1\, \mathrm{d}M_2. \tag{1-1}$$

The parameter N is a scaling parameter; when considering the asymptotic regime $n \to \infty$ the parameter N is also sent to infinity in such a way that $\frac{n}{N} \to T \in \mathbb{R}_+$. We will assume henceforth $T = 1$ and that $N = n$.

It was shown in [2] that, similar to other integrable matrix models, all correlation functions of the eigenvalues of the two matrices M_1, M_2 can be computed exactly in terms of certain *Cauchy biorthogonal polynomials* (BOPs). The latter consist of two sequences of polynomials $\{p_n(x), q_n(y)\}_{n\in\mathbb{N}}$ of exact degree n with the defining properties

$$\int_{\mathbb{R}_+^2} \frac{\mathrm{e}^{-N(V_1(x)+V_2(y))}}{x+y} p_\ell(x) q_m(y) \mathrm{d}x\, \mathrm{d}y = \delta_{\ell m}\ , \quad p_n(x) = c_n x^n + \dots, \quad q_n(y) = c_n y^n + \dots . \quad c_n > 0. \tag{1-2}$$

In [3] the algebraic properties of these polynomials were investigated but no concrete example which could be considered "classical" was provided. On the other hand, even before the Cauchy BOPs were introduced, an instance of such polynomials associated with a classical weight appeared implicitly in [4] in the study of the biorthogonal Laguerre ensemble, one of several examples of biorthogonal ensembles considered there that allow an explicit computation of correlation functions. In this paper, we apply the machinery developed for general Cauchy BOPs in [2],[3] to this particular instance. More precisely, we shall consider the case $V_1(x) = x + \frac{a}{N}\ln x,\ V_2(y) = y + \frac{b}{N}\ln y$; in this case the scaling factor N in eq. (1-2) can be absorbed in a common rescaling of the variables x, y without affecting the general properties by setting $V_1(x) = \frac{1}{N}(x + a\ln x),\ V_2(y) = \frac{1}{N}(y + b\ln y)$. The corresponding biorthogonal polynomials are thus defined by

$$\int_{\mathbb{R}_+^2} \frac{x^a y^b \mathrm{e}^{-x-y}}{x+y} p_\ell(x) q_m(y) = \delta_{\ell m}\ , \quad p_n(x) = c_n x^n + \dots, \quad q_n(y) = c_n y^n + \dots . \quad c_n > 0. \tag{1-3}$$

The present paper has three main goals:

1. obtain explicit formulæ for p_n, q_n in eq. (1-3) and for related objects;
2. find explicit formulæ for the correlation functions at finite n;

3. formulate a scaling limit of the correlation functions near the origin and thus define a limiting random point field.

We shall show that all the objects naturally appearing in the investigation of the BOPs defined by eq. (1-3) are intimately related to Meijer-G functions. It is for this reason that we shall name the limiting process mentioned in the last item above the *Meijer-G process*.

The results of the paper are summarized in the next section, while the remainder of the paper is devoted to the proof of the theorems stated in Section 1.1.

1.1 Summary of results

We recall the results of [2] (collected and explained in Appendix B, in particular eq. (B.2b)). The correlation functions of the Cauchy two-matrix model are expressed as determinants

$$R^{(k,\ell)}(x_1,\dots,x_k;y_1,\dots,y_\ell)=\det\left[\begin{array}{c|c}\underset{1\le i,j\le k}{H_{01}^{(n)}(x_i,x_j)} & \underset{1\le i\le k;1\le j\le \ell}{H_{00}^{(n)}(x_i,y_j)}\\ \hline \underset{1\le i\le \ell;1\le j\le k}{H_{11}^{(n)}(y_i,x_j)} & \underset{1\le i,j\le \ell}{H_{10}^{(n)}(y_i,y_j)}\end{array}\right], \tag{1-4}$$

where the kernels $H_{\rho\mu}^{(n)}$ are given by

$$H_{00}^{(n)}(x,y):=x^a y^b \mathrm{e}^{-x-y}K_{00}^{(n)}(x,y), \qquad H_{01}^{(n)}(x,y):=x^a\mathrm{e}^{-x}K_{01}^{(n)}(x,y), \tag{1-5a}$$

$$H_{10}^{(n)}(x,y):=y^b\mathrm{e}^{-y}K_{10}^{(n)}(x,y), \qquad H_{11}^{(n)}(x,y):=K_{11}^{(n)}(x,y), \tag{1-5b}$$

with the kernels $K_{\mu\nu}^{(n)}$ defined in terms of the Cauchy biorthogonal polynomials p_n,q_n as:

$$K_{00}^{(n)}(x,y):=\sum_{j=0}^{n-1}p_j(x)q_j(y), \qquad K_{01}^{(n)}(x,y):=\int_{\mathbb{R}_+}K_{00}^{(n)}(x,y')\frac{y'^b\mathrm{e}^{-y'}\mathrm{d}y'}{y+y'}, \tag{1-6a}$$

$$K_{10}^{(n)}(x,y):=\int_{\mathbb{R}_+}K_{00}^{(n)}(x',y)\frac{x'^a\mathrm{e}^{-x'}\mathrm{d}x}{x+x'} \qquad K_{11}^{(n)}(x,y):=\int_{\mathbb{R}_+^2}K_{00}^{(n)}(x',y')\frac{x'^a\mathrm{e}^{-x'}\mathrm{d}x'}{x+x'}\frac{y'^b\mathrm{e}^{-y'}\mathrm{d}y'}{y+y'}-\frac{1}{x+y}. \tag{1-6b}$$

Our first main result is a compact expression for these four kernels for finite n; to present it we set $\alpha=a+b$ and define four functions:

$$G_{c,n}(z):=\int_\gamma\frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(u)\Gamma(n+\alpha+1-u)}{\Gamma(c+1-u)\Gamma(n+u)\Gamma(\alpha-u+1)}z^{-u}, \tag{1-7}$$

$$\widetilde{G}_{c,n}(w):=\int_\gamma\frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(u-c)\Gamma(u)\Gamma\left(n+\alpha+1-u\right)}{\Gamma(\alpha+1-u)\Gamma\left(n+u\right)}w^{-u}, \tag{1-8}$$

and

$$H_{c,n}(z) := \int_{\gamma+c} \frac{\mathrm{d}u}{2\pi i} \frac{\Gamma(u+c)\Gamma(n+\alpha-c+1-u)}{\Gamma(1-u)\Gamma(c+n+u)\Gamma(\alpha-c-u+1)} z^{-u} = z^c G_{c,n}(z), \tag{1-9}$$

$$\widetilde{H}_{c,n}(w) := \int_{\gamma+c} \frac{\mathrm{d}u}{2\pi i} \frac{\Gamma(u)\Gamma(u+c)\Gamma\left(n+\alpha-c+1-u\right)}{\Gamma(\alpha-c+1-u)\Gamma\left(c+n+u\right)} w^{-u} = w^c G_{c,n}(w). \tag{1-10}$$

In the above expressions γ is a contour originating at $-\infty$ in the lower half-plane and returning to $-\infty$ in the upper half-plane in such a way as to leave all the poles of the Γ functions in the numerator containing the variable $+u$ inside the contour, while leaving those of the Γ functions of variable $-u$ outside. Such types of integrals are Mellin-Barnes integrals and the expressions above are special cases of Meijer-G functions (see [5], 5.3, p. 206). Then

Theorem 1.1. *The kernels $H^{(n)}_{\mu\nu}, K^{(n)}_{\mu\nu}$ are given by*

$$\begin{aligned}
&H^{(n)}_{00}(x,y) = \int_0^1 H_{a,n}(tx)H_{b,n}(ty)\mathrm{d}t, && H^{(n)}_{01}(x,y) = \int_0^1 H_{a,n}(tx)\widetilde{H}_{a,n}(ty)\mathrm{d}t,\\
&H^{(n)}_{10}(x,y) = \int_0^1 \widetilde{H}_{a,n}(tx)H_{b,n}(ty)\mathrm{d}t, && H^{(n)}_{11}(x,y) = \int_0^1 \widetilde{H}_{a,n}(tx)\widetilde{H}_{b,n}(ty)\mathrm{d}t - \frac{1}{x+y},\\
&K^{(n)}_{00}(x,y) = \int_0^1 G_{a,n}(tx)G_{b,n}(ty)t^\alpha\mathrm{d}t, && K^{(n)}_{01}(x,y) = y^b\mathrm{e}^y\int_0^1 G_{a,n}(tx)\widetilde{G}_{b,n}(ty)t^\alpha\mathrm{d}t,\\
&K^{(n)}_{10}(x,y) = x^a\mathrm{e}^x\int_0^1 \widetilde{G}_{a,n}(tx)G_{b,n}(ty)t^\alpha\mathrm{d}t, && K^{(n)}_{11}(x,y) = x^by^a\mathrm{e}^{x+y}\int_0^1 \widetilde{G}_{a,n}(tx)\widetilde{G}_{b,n}(ty)t^\alpha\mathrm{d}t - \frac{\mathrm{e}^{x+y}}{x+y}.
\end{aligned}$$

The above theorem is the summary of Theorems 3.5,3.6,3.7 that contain the computation of the kernels $K^{(n)}_{\mu\nu}$, whereas the ensuing expressions for the kernels $H^{(n)}_{\mu\nu}$ is a simple rewrite using the definitions (eq. (1-5), 1-6, 1-9, and eq. (1-10)).

1.1.1 The scaling random field at the origin: the Meijer-G random point field

With the substitutions $x = \zeta n^{-2}, y = \xi n^{-2}$ we arrive in the limit $n \to \infty$ at a novel universality class of random point fields that we name the *Meijer-G random point field.* First we define

Definition 1.1. *Let*

$$H_c(\zeta) := \zeta^c G_c(\zeta) = \int_\gamma \frac{\mathrm{d}u}{2i\pi} \frac{\Gamma(u+c)}{\Gamma(1-u)\Gamma(\alpha-c+1-u)} \zeta^{-u}, \tag{1-11}$$

$$\widetilde{H}_c(\zeta) := \zeta^c \widetilde{G}_c(\zeta) = \int_\gamma \frac{\mathrm{d}u}{2i\pi} \frac{\Gamma(u)\Gamma(u+c)}{\Gamma(\alpha-c+1-u)} \zeta^{-u}. \tag{1-12}$$

The contour γ is a contour of the form in Fig. 1 linking to all the poles in the numerators of the integrands. Note that $H_c, \widetilde{H}_c$ are Meijer-G functions as in [5] and 16.17 in DLMF [1] (see definition in App. A).

---

[1] DLMF="Digital Library of Mathematical Functions", http://dlmf.nist.gov

Theorem 1.2 (Meijer-G random point field and universal class)**.** *In the scaling limit the correlations of the eigenvalues of M_1, M_2 are determined by the two–level random point field with kernels below*

$$\lim_{n\to\infty} n^{-2} H_{00}^{(n)}\left(\frac{\zeta}{n^2}, \frac{\xi}{n^2}\right) = \mathcal{G}_{00}(\zeta,\xi), \qquad \lim_{n\to\infty} n^{-2} H_{01}^{(n)}\left(\frac{\zeta}{n^2}, \frac{\xi}{n^2}\right) = \mathcal{G}_{01}(\zeta,\xi), \tag{1-13}$$

$$\lim_{n\to\infty} n^{-2} H_{10}^{(n)}\left(\frac{\zeta}{n^2}, \frac{\xi}{n^2}\right) = \mathcal{G}_{10}(\zeta,\xi), \qquad \lim_{n\to\infty} n^{-2} H_{11}^{(n)}\left(\frac{\zeta}{n^2}, \frac{\xi}{n^2}\right) = \mathcal{G}_{11}(\zeta,\xi), \tag{1-14}$$

where

$$\mathcal{G}_{00}(\zeta,\xi) = \int_0^1 H_a(t\zeta) H_b(t\xi) \mathrm{d}t, \qquad \mathcal{G}_{01}(\zeta,\xi) = \int_0^1 H_a(t\zeta) \widetilde{H}_b(t\xi) \mathrm{d}t, \tag{1-15}$$

$$\mathcal{G}_{10}(\zeta,\xi) = \int_0^1 \widetilde{H}_a(t\zeta) H_b(t\xi) \mathrm{d}t \qquad \mathcal{G}_{11}(\zeta,\xi) = \int_0^1 \widetilde{H}_a(t\zeta) \widetilde{H}_b(t\xi) \mathrm{d}t - \frac{1}{\zeta+\xi}, \tag{1-16}$$

with $H_c, \widetilde{H}_c$ as in Definition 1.1. The convergence is uniform for ξ, η within compact sets and the error approximation is within $\mathcal{O}(n^{-1})$.

Proposition 3.2 provides alternative expressions for the kernels $\mathcal{G}_{\mu\nu}$ in terms of "point-split bilinear concomitants". Section 3.3 is devoted to the proof of Theorem 1.2. We expect the following conjecture to be true.

Conjecture 1.1. *The Meijer random process obtained in the scaling limit in this paper is universal within the class of Cauchy matrix models of the form*

$$\mathrm{d}\mu(M_1, M_2) = \frac{1}{\mathcal{Z}_n} \det(M_1)^a \det(M_2)^b \frac{\mathrm{e}^{-n\mathrm{Tr}(V_1(M_1)+V_2(M_2))}}{\det(M_1+M_2)^n} \mathrm{d}M_1 \mathrm{d}M_2$$

with V_i analytic near the origin and the scaling $x \mapsto x\, n^{-3}$.

1.1.2 Outlook

The issue as to the real nature of the universality of the Meijer-G random point field for arbitrary potentials will be addressed in a forthcoming publication; the proof will be based on the nonlinear steepest descent method. An interesting problem which will also be addressed in a separate publication is to connect the gap probability of the Meijer-G field to a suitable Painlevé transcendent, namely, to obtain the analog of the Tracy–Widom distribution of the largest eigenvalue [12].

Acknowledgements This paper was initiated at the Banff International Research Station. We thank BIRS for the hospitality and for providing excellent work conditions. M.B. wishes to thank SISSA (Trieste) for hospitality during which part of the work was completed. M. B. and J.S. acknowledge a support by Natural Sciences and Engineering Research Council of Canada. M.G. is partially supported by the NSF grant DMS-1101462.

2 Setup

As was observed in [4] the Cauchy biorthogonal polynomials in eq. (1-3) are related to the classical Jacobi orthogonal polynomials for the weight $x^{a+b}\mathrm{d}x$ on $[0,1]$. We thank A. Borodin for pointing out this connection whose main point is as follows: consider the bi-moment matrix

$$I_{ij} = \iint_0^{+\infty} x^i y^j \frac{e^{-(x+y)}}{x+y} x^a y^b \, d\, x d\, y \ .$$

With the change of variables $x = rs$, $y = r(1-s)$ the integral becomes

$$\int_0^{+\infty} r^{i+j+a+b} e^{-r} \mathrm{d}r \int_0^1 \frac{t^{i+a+1}(1-t)^{j+b+1}\mathrm{d}t}{t(1-t)} = \frac{\Gamma(a+i+1)\Gamma(b+j+1)}{(i+a+j+b+1)} \ . \tag{2-1}$$

Notice now that the Hankel moment matrix M for the Jacobi polynomials on $[0,1]$ with weight x^{a+b} is given by

$$M_{ij} = \int_0^1 x^{i+j} x^{a+b} d\, x = \frac{1}{i+j+a+b+1} \Rightarrow I_{ij} = \Gamma(a+i+1) M_{ij} \Gamma(b+j+1). \tag{2-2}$$

This immediately implies the following Proposition.

Proposition 2.1. *Let* $P_m(x) = \sum_{i=1}^m C_{m,i} x^i$ *denote the is the* m*-th Jacobi orthogonal polynomial*[2] *then*

$$\widetilde{p}_m(x) := \sum_{i=0}^m C_{m,i} \frac{x^i}{\Gamma(a+i+1)}, \qquad \widetilde{q}_n(y) := \sum_{j=1}^n C_{n,j} \frac{y^j}{\Gamma(b+j+1)} \tag{2-3}$$

are the Cauchy biorthogonal polynomials (not orthonormal) associated with densities $x^a \mathrm{e}^{-x}$ *and* $y^b \mathrm{e}^{-y}$ *and*

$$\int_{\mathbb{R}_+^2} \widetilde{p}_m(x) \widetilde{q}_\ell(y) \frac{x^a y^b \mathrm{e}^{-x-y}}{x+y} \mathrm{d}x\mathrm{d}y = \int_0^1 P_m(x) P_\ell(x) x^\alpha \mathrm{d}x = h_m \delta_{m\ell} \tag{2-4}$$

where $h_m = \frac{1}{2m+\alpha+1}$. *This result is valid as long as* $a+1>0, b+1>0, a+b+1>0$.

Remark 2.1. The partition function of the corresponding Cauchy two-matrix model can be easily computed using Proposition 2.1 : in [2] it was shown that if c_n are the norms of the *monic* polynomials (denoted here provisionally by $p_n^{(monic)}, q_n^{(monic)}$)

$$\int_{\mathbb{R}_+^2} p_m^{(monic)}(x) q_\ell^{(monic)}(y) \frac{x^a y^b \mathrm{e}^{-x-y}}{x+y} \mathrm{d}x\mathrm{d}y = c_m \delta_{m\ell} \tag{2-5}$$

---

[2] Here we define the orthogonality on $[0,1]$ rather than on $[-1,1]$. The latter is more customary, but the correspondence between the two versions is a simple affine transformation of the independent variable $x \to 2x-1$.

then $\mathcal{Z}_n = (n!)^2 \prod_{j=0}^{n-1} c_j$. Inspection of the leading coefficients of $\widetilde{p}_n, \widetilde{q}_n$ in Theorem 3.1 shows that

$$c_n = \left[\frac{n!\Gamma(\alpha+n+1)}{\Gamma(\alpha+2n+1)}\right]^2 \frac{\Gamma(a+n+1)\Gamma(b+n+1)}{2n+\alpha+1} = \tag{2-6}$$

$$= \left[\frac{n!\Gamma(\alpha+n+1)}{\Gamma(n+\frac{\alpha+1}{2})\Gamma(n+1+\frac{\alpha}{2})}\right]^2 \frac{\Gamma(a+n+1)\Gamma(b+n+1)\pi}{2^{2n+\alpha}(n+\frac{\alpha+1}{2})} \tag{2-7}$$

$$\mathcal{Z}_n = (n!)^2 \prod_{j=0}^{n-1} c_j = \frac{\Gamma\left(\frac{\alpha+1}{2}\right)}{\Gamma\left(n+\frac{\alpha+1}{2}\right)} \frac{G(n+1)^2 G(\alpha+n+1)^2 G(\frac{\alpha+1}{2})^2 G(\frac{\alpha}{2}+1)^2}{G(\alpha+1)^2 G(\frac{\alpha+1}{2}+n)^2 G(n+1+\frac{\alpha}{2})^2} \frac{G(a+n+1)G(b+n+1)\pi}{G(a+1)G(b+1)2^{n(n-1)+n\alpha}}\,. \tag{2-8}$$

Here (and only in the above formula) $G(z)$ denotes Barnes' G-function, satisfying the relation $G(z+1) = \Gamma(z)G(z)\ ,\quad G(1) = 1$.

2.1 The kernels of the correlation functions

The statistics of eigenvalues of M_1, M_2 is expressible in terms of four kernels that can be expressed in terms of the CBOPs and auxiliary functions. In keeping with the notation of [3] we introduce the auxiliary functions

$$p_n^{(1)}(z) := \int_{\mathbb{R}_+} \frac{p_n(x) x^a \mathrm{e}^{-x} \mathrm{d}x}{z-x}\ , \qquad q_n^{(1)}(w) := \int_{\mathbb{R}_+} \frac{q_n(y) y^a \mathrm{e}^{-y} \mathrm{d}y}{w-y}. \tag{2-9}$$

The four kernels to be computed are

$$K_{00}^{(n)}(x,y) := \sum_{j=0}^{n-1} p_j(x) q_j(y), \tag{2-10}$$

$$K_{01}^{(n)}(x,y) := -\sum_{j=0}^{n-1} p_j(x) q_j^{(1)}(-y) = \int_{\mathbb{R}_+} \frac{y'^b \mathrm{e}^{-y'} \mathrm{d}y}{y+y'} K_{00}^{(n)}(x,y'), \tag{2-11}$$

$$K_{10}^{(n)}(x,y) := -\sum_{j=0}^{n-1} p_j^{(1)}(-x) q_j(y) = \int_{\mathbb{R}_+} \frac{x'^a \mathrm{e}^{-x'} \mathrm{d}x}{x+x'} K_{00}^{(n)}(x',y), \tag{2-12}$$

$$K_{11}^{(n)}(x,y) := \sum_{j=0}^{n-1} p_j^{(1)}(-x) q_j^{(1)}(-y) - \frac{1}{x+y} = \tag{2-13}$$

$$= \int_{\mathbb{R}_+^2} K_{00}^{(n)}(x',y') \frac{x'^a \mathrm{e}^{-x'} \mathrm{d}x'}{x+x'} \frac{y'^b \mathrm{e}^{-y'} \mathrm{d}y'}{y+y'} - \frac{1}{x+y}. \tag{2-14}$$

Each of these kernels can be recovered in general by solving a Riemann–Hilbert problem [3, 2] but in the present setting this will be a direct computation.

2.2 Facts on Jacobi Polynomials

Since Jacobi polynomials will be instrumental to our computations, we recall their basic properties. Jacobi polynomials are defined by the formulæ

$$P_n^{(\alpha,\beta)}(\xi) = \frac{\Gamma(\alpha+n+1)}{n!\,\Gamma(\alpha+\beta+n+1)} \sum_{m=0}^{n} \binom{n}{m} \frac{\Gamma(\alpha+\beta+n+m+1)}{\Gamma(\alpha+m+1)} \left(\frac{\xi-1}{2}\right)^m ,$$

$$\int_{-1}^{1} (1-\xi)^\alpha (1+\xi)^\beta P_m^{(\alpha,\beta)}(\xi) P_n^{(\alpha,\beta)}(\xi)\, d\xi = \frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)n!} \delta_{nm}.$$

We shall use the variable $z = \frac{1-\xi}{2}$ so that the orthogonality becomes

$$\int_0^1 z^\alpha (1-z)^\beta P_m^{(\alpha,\beta)}(1-2z)\, P_n^{(\alpha,\beta)}(1-2z)\, dz = \frac{1}{2n+\alpha+\beta+1} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)n!} \delta_{nm}. \tag{2-15}$$

The case of interest to us is $\beta = 0, \ \alpha = a+b$ and thus we shall introduce a simplified notation

$$P_n(z) := P_n^{(\alpha,0)}(1-2z) = \sum_{m=0}^{n} \frac{\Gamma(\alpha+n+m+1)}{m!(n-m)!\Gamma(\alpha+m+1)} (-z)^m \tag{2-16}$$

$$= \int_\gamma \frac{\Gamma(\alpha+n+1-u)\Gamma(u)}{\Gamma(n+1+u)\Gamma(\alpha-u+1)} z^{-u} \frac{\mathrm{d}u}{2i\pi} = \frac{\Gamma(n+\alpha+1)}{n!\Gamma(\alpha+1)} {}_2F_1\left(\begin{matrix} -n, n+\alpha+1 \\ \alpha+1 \end{matrix} ; z\right), \tag{2-17}$$

$$\int_0^1 P_m(z)\, P_n(z) z^\alpha\, dz = \frac{1}{2n+\alpha+1} \delta_{nm}, \tag{2-18}$$

where the integral representation is valid for $0 < |z| < 1$. In eq. (2-17) the contour embraces $u \in \mathbb{R}_-$ (Fig. 1); note that the poles of $\Gamma(u)$ for $u = -n-1, -n-2, \dots$ are cancelled by the poles of the term $\Gamma(n+1+u)$ in the denominator and thus the result is a polynomial as a consequence of a simple residue computation.

The Christoffel-Darboux kernel for Jacobi polynomials [10] is defined as

$$J_n(z,w) := \sum_{m=0}^{n-1} \frac{P_m(z)P_m(w)}{h_m} \tag{2-19}$$

where h_n are the norms squared of P_n in eq. (2-18). The following proposition is simple but we could not find it in the literature, and since its proof is very close to proofs later on we present it here.



Figure 1: The typical contour for Meijer-type integrals.

Proposition 2.2. *For $\beta = 0$, the Christoffel-Darboux kernel for Jacobi polynomials can be expressed as*

$$K_n(z,w) := \sum_{j=0}^{n-1} (2j+\alpha+1) P_j(z) P_j(w) =$$

$$= \int_{\gamma^2} \frac{\mathrm{d}u}{2\pi i} \frac{\mathrm{d}v}{2\pi i} \frac{z^{-u} w^{-v}}{1+\alpha-v-u} \frac{\Gamma(u)\Gamma(v)\Gamma(n+\alpha+1-u)\Gamma(n+\alpha+1-v)}{\Gamma(n+u)\Gamma(n+v)\Gamma(\alpha+1-u)\Gamma(\alpha+1-v)}, \tag{2-20}$$

valid for $0 < |z|, |w| < 1$. *Alternatively, introducing the function* $\widehat{G}_n(z)$ (a polynomial of degree $n-1$)

$$\widehat{G}_n(z) := \int_\gamma \frac{\mathrm{d}u}{2\pi i} \frac{\Gamma(u)\Gamma(n+\alpha+1-u)}{\Gamma(n+u)\Gamma(\alpha+1-u)} z^{-u} \tag{2-21}$$

we can write

$$K_n(z,w) = \int_0^1 \widehat{G}_n(tz)\widehat{G}_n(tw)t^\alpha \mathrm{d}t. \tag{2-22}$$

At this point we recall the definition of the Meijer G functions, or rather a class of G-functions pertinent to this paper.

Definition 2.1 (Meijer G-functions)**.** *Suppose two pairs of natural numbers* $p \le q$ *and* $0 \le m \le q$, $0 \le n \le p$ *are given. Then the Meijer* G*-function is defined by its Mellin-Barnes integral as follows:*

$$G^{m,n}_{p,q}\left(\begin{array}{c} a_1, \dots, a_p \\ b_1, \dots, b_q \end{array} \middle| \, z \right) = \int_\gamma \frac{\mathrm{d}u}{2\pi i} \frac{\prod_{j=1}^m \Gamma(b_j+u) \prod_{j=1}^n \Gamma(1-a_j-u)}{\prod_{j=m+1}^q \Gamma(1-b_j-u) \prod_{j=n+1}^p \Gamma(a_j+u)} z^{-u}, \tag{2-23}$$

The contour γ*, depicted in Fig. 1, is chosen to encircle all the poles of functions* $\Gamma(b_j+u)$ *and none of the poles of functions* $\Gamma(1-a_j-u)$ *(the implicit assumption is that none of the poles of the former coincides with any of the poles of the latter).*

Further properties of G-functions are discussed in the Appendix A. We can identify all the functions discussed so far in terms of G-functions. We have

Proposition 2.3. *The Jacobi polynomials* P_n *and the function* $\widehat{G}_n$ *appearing in the representation of the Christoffel-Darboux kernel are* G*-functions with symbols*

$$P_n(z) = G^{1,1}_{2,2}\left(\begin{array}{c} -\alpha-n, n+1 \\ 0, -\alpha \end{array} \middle| \, z \right), \quad \widehat{G}_n(z) = G^{1,1}_{2,2}\left(\begin{array}{c} -\alpha-n, n \\ 0, -\alpha \end{array} \middle| \, z \right).$$

We shall need the following Lemma, whose elementary proof is omitted.

Lemma 2.1. *Let* $\Omega^j_k = \frac{\Gamma(\alpha+j+1-u)\Gamma(\alpha+j+1-v)}{\Gamma(k+u)\Gamma(k+v)}$. *Then*

$$\sum_{j=0}^{n-1} (2j+\alpha+1)\Omega^j_{j+1} = \frac{\Omega^n_n - \Omega^0_0}{1+\alpha-u-v} = \int_0^1 t^{\alpha-u-v}(\Omega^n_n - \Omega^0_0)\mathrm{d}t.$$

Proof. [**Proof of Prop. 2.2**] From eq. (2-19) and the value of the norms squared given by eq. (2-18) we have

$$\sum_{j=0}^{n-1}(2j+\alpha+1)P_n(z)P_n(w) =$$

$$\int_{\gamma\times\gamma} \frac{\mathrm{d}u}{2\pi i}\frac{\mathrm{d}v}{2\pi i} \frac{z^{-u}w^{-v}\Gamma(u)\Gamma(v)}{\Gamma(\alpha-u+1)\Gamma(\alpha-v+1)} \sum_{j=0}^{n-1} \frac{(2j+\alpha+1)\Gamma(\alpha+j+1-u)\Gamma(\alpha+j+1-v)}{\Gamma(j+1+u)\Gamma(j+1+v)}. \tag{2-24}$$

Applying now Lemma 2.1 to eq. (2-24) we obtain

$$\int_{\gamma\times\gamma} \frac{\mathrm{d}u}{2\pi i}\frac{\mathrm{d}v}{2\pi i}\frac{z^{-u}w^{-v}}{1+\alpha-u-v}\left[\frac{\Gamma(u)\Gamma(v)\Gamma(n+\alpha+1-u)\Gamma(n+\alpha+1-v)}{\Gamma(n+u)\Gamma(n+v)\Gamma(\alpha-u+1)\Gamma(\alpha-v+1)}-1\right]=$$
$$\int_{\gamma\times\gamma} \frac{\mathrm{d}u}{2\pi i}\frac{\mathrm{d}v}{2\pi i}\int_0^1 t^{\alpha}(tz)^{-u}(tw)^{-v}\left[\frac{\Gamma(u)\Gamma(v)\Gamma(n+\alpha+1-u)\Gamma(n+\alpha+1-v)}{\Gamma(n+u)\Gamma(n+v)\Gamma(\alpha-u+1)\Gamma(\alpha-v+1)}-1\right]\mathrm{d}t.$$

The term with the 1 vanishes identically because the contours of integration can be retracted to $-\infty$ and $z, w \in (0,1)$, and the denominator does not vanish ($\alpha > -1$) because the contour around 0 can be moved sufficiently to the left so that the denominator has no zero anywhere inside the contour. This immediately yields the first formula eq. (2-20). To obtain eq. (2-22) we switch the order of integration in the second integral formula above. □

It is interesting to consider the scaling limit in which $n \to \infty$ and x, y are rescaled by n^{-2}; using the Stirling approximation formula one obtains readily the following Lemma.

Lemma 2.2. *We have the following asymptotic formula*

$$\frac{\Gamma(z+\delta)}{\Gamma(z+\rho)} = z^{\delta-\rho}\left(1+\frac{(\delta+\rho-1)(\delta-\rho)}{2z}+\mathcal{O}(z^{-2})\right), \qquad |\arg(z)|<\pi, \tag{2-25}$$

where the estimate is uniform in each proper subsector.

Then, using Lemma 2.2 and Proposition 2.2

$$\begin{aligned}
&K_n(xn^{-2}, yn^{-2}) = \\
&= n^{2\alpha+2}\int_{\gamma^2}\frac{\mathrm{d}u}{2i\pi}\frac{\mathrm{d}v}{2i\pi}\frac{x^{-u}y^{-v}}{1+\alpha-v-u}\frac{\Gamma(u)\Gamma(v)}{\Gamma(\alpha-u+1)\Gamma(\alpha-v+1)}\left(1+\frac{\alpha(\alpha+1-u-v)}{n}+\mathcal{O}(n^{-2})\right) = \\
&= n^{2\alpha+2}\left(1+\frac{\alpha(\alpha+1+x\partial_x+y\partial_y)}{n}+\mathcal{O}(n^{-2})\right)\int_{\gamma^2}\frac{\mathrm{d}u}{2i\pi}\frac{\mathrm{d}v}{2i\pi}\int_0^1\frac{\Gamma(u)\Gamma(v)t^{\alpha}(tx)^{-u}(ty)^{-v}\mathrm{d}t}{\Gamma(\alpha-u+1)\Gamma(\alpha-v+1)} = \\
&= n^{2\alpha+2}\left(1+\frac{\alpha(\alpha+1+x\partial_x+y\partial_y)}{n}+\mathcal{O}(n^{-2})\right)\int_0^1 t^{\alpha}B_\alpha(tx)B_\alpha(ty)\mathrm{d}t,
\end{aligned} \tag{2-26}$$

where we have introduced the function B_α defined as

$$B_\alpha(z) := \int_\gamma \frac{\mathrm{d}u}{2i\pi}z^{-u}\frac{\Gamma(u)}{\Gamma(\alpha-u+1)} = z^{-\alpha/2}J_\alpha(2\sqrt{z}). \tag{2-27}$$

In the above J_α denotes the Bessel function of the first kind. We also note the integral makes sense for any value of z thanks to the super-exponential behavior of the gamma functions. The leading term of the kernel in eq. (2-26) is almost the same as the Bessel kernel. Indeed, using eq. (2.2) in [11] we can write the Bessel kernel, denoted here by K_B, as

$$K_B(x,y) := \frac{J_\alpha(\sqrt{x})\sqrt{y}J_\alpha'(\sqrt{y}) - J_\alpha(\sqrt{y})\sqrt{x}J_\alpha'(\sqrt{x})}{2(x-y)} = \frac{1}{4}\int_0^1 J_\alpha(\sqrt{xt})J_\alpha(\sqrt{yt})\mathrm{d}t. \tag{2-28}$$

A direct inspection shows that

$$K_{_B}(4x,4y) = \frac{1}{4}\int_0^1 J_\alpha(2\sqrt{xt})J_\alpha(2\sqrt{yt})\mathrm{d}t = (xy)^{\frac{\alpha}{2}}\frac{1}{4}\int_0^1 t^\alpha B_\alpha(tx)B_\alpha(ty)\mathrm{d}t. \tag{2-29}$$

Thus

$$K_n(xn^{-2}, yn^{-2}) = n^{2\alpha+2}\left(1 + \frac{\alpha(\alpha+1+x\partial_x+y\partial_y)}{n} + \mathcal{O}(n^{-2})\right)\frac{4K_B(4x,4y)}{(xy)^{\frac{\alpha}{2}}}$$

This is not surprising since the Bessel kernel arises precisely as the scaling limit of the Jacobi matrix model near the hard-edge.

3 Cauchy-Laguerre Biorthogonal polynomials

The reason for the use of Laguerre's name is clearly justified by the shape of the weights; nevertheless, as we are going to see, the resulting polynomials are more closely related to Jacobi polynomials. Using the explicit form of Jacobi polynomials as given by eq. (2-16) and their relation to Cauchy BOPs (see eq. (2-3)) we can immediately assert the following

Theorem 3.1. *The polynomials $\widetilde{p}_n$ of Proposition 2.1 are given by*

$$\widetilde{p}_n(z) = \sum_{j=0}^{n}\frac{\Gamma(\alpha+n+j+1)(-z)^j}{j!(n-j)!\Gamma(\alpha+j+1)\Gamma(a+j+1)} = \frac{\Gamma(\alpha+n+1)}{n!\Gamma(\alpha+1)\Gamma(a+1)}\,{}_2F_2\left(\begin{matrix}-n, \alpha+n+1\\ a+1, \alpha+1\end{matrix}; z\right) =$$

$$= \int_\gamma \frac{\mathrm{d}u}{2i\pi}\frac{\Gamma(\alpha+n-u+1)\Gamma(u)}{\Gamma(n+1+u)\Gamma(\alpha+1-u)\Gamma(a+1-u)}z^{-u} \tag{3-1}$$

where the contour γ is the same as in Fig. 1 with the poles of $\Gamma(\alpha+n+1-u)$ in its exterior. The integral representation is valid for any $z \neq 0$. The expressions for the polynomials $\widetilde{q}_n$ follow from those for $\widetilde{p}_n$ by exchanging $a \leftrightarrow b$.

Proof. The expression in terms of (generalized) hypergeometric function follows from its definition (see 16.2 in DLMF); he integral representation is a simple consequence of the Cauchy residue theorem and the fact that $\mathop{\mathrm{res}}_{u=-k}\Gamma(u) = (-1)^k\frac{1}{k!}$. □

We have two simple corollaries

Corollary 3.1 (Averages of $\widetilde{p}_n, \widetilde{q}_n$)**.** *The average of $\widetilde{p}_n$ with respect to the measure $x^a\mathrm{e}^{-x}\mathrm{d}x$, ($\widetilde{q}_n$* with respect to $y^b\mathrm{e}^{-y}\mathrm{d}y$) is

$$\int_0^\infty x^a\mathrm{e}^{-x}\widetilde{p}_n(x)\mathrm{d}x = (-1)^n = \int_0^\infty y^a\mathrm{e}^{-y}\widetilde{q}_n(y)\mathrm{d}y. \tag{3-2}$$

Proof. A simple computation gives:

$$\int_0^\infty x^a\mathrm{e}^{-x}\widetilde{p}_n(x)\mathrm{d}x = \sum_{j=0}^{n}\frac{\Gamma(\alpha+n+j+1)(-1)^j\int_0^\infty x^{j+a}\mathrm{e}^{-x}\mathrm{d}x}{j!(n-j)!\Gamma(\alpha+j+1)\Gamma(a+j+1)} = \sum_{j=0}^{n}\frac{\Gamma(\alpha+n+j+1)(-1)^j}{j!(n-j)!\Gamma(\alpha+j+1)} = P_n(1).$$

It is well known that $P_n(1) = (-1)^n$ as a consequence of eq. (2-16). **Q.E.D.**

The second corollary is an elementary identification of $\widetilde{p}_n$, $\widetilde{q}_n$ respectively, with one of the Meijer G-functions. As we will demonstrate the Meijer G-functions play a significant role in the forthcoming computations.

Corollary 3.2. *The polynomials $\widetilde{p}_n$ and $\widetilde{q}_n$ can be identified with the following Meijer G-functions:*

$$\widetilde{p}_n(z) = G^{1,1}_{2,3}\left(\begin{matrix} -\alpha - n, n+1 \\ 0, -\alpha, -a \end{matrix} \middle| z \right), \qquad \widetilde{q}_n(z) = G^{1,1}_{2,3}\left(\begin{matrix} -\alpha - n, n+1 \\ 0, -\alpha, -b \end{matrix} \middle| z \right).$$

Direct inspection of the leading coefficient of $\widetilde{p}_n, \widetilde{q}_n$ together with Corollary 3.1 imply the next result

Theorem 3.2. *Let $\pi_n := \sqrt{2n+\alpha+1}\sqrt{\frac{\Gamma(a+n+1)}{\Gamma(b+n+1)}}$, $\eta_n := \sqrt{2n+\alpha+1}\sqrt{\frac{\Gamma(b+n+1)}{\Gamma(a+n+1)}}$. Then the* **bi-orthonormal polynomials** *p_n, q_n, normalized to have identical positive leading coefficients, are given by*

$$p_n(z) = (-1)^n \pi_n \widetilde{p}_n, \quad q_n(z) = (-1)^n \eta_n \widetilde{q}_n.$$

Moreover, $\pi_n = \int_0^\infty x^a \mathrm{e}^{-x} p_n(x) \mathrm{d}x$ and $\eta_n = \int_0^\infty y^b \mathrm{e}^{-y} q_n(y) \mathrm{d}y$.

3.1 The auxiliary functions $p_n^{(1)}$ and $p_n^{(2)}$

Since the definition of $q_n^{(1)}$ can be obtained from that of $p_n^{(1)}$ by swapping a with b we subsequently will focus only on $p_n^{(1)}$ as defined in eq. (2-9).

Our goal is to express $p_n^{(1)}(z)$ in terms of the G-functions. To this end we first compute $\int_0^\infty \frac{x^a \mathrm{e}^{-x} \widetilde{p}_n(x)}{z+x} \mathrm{d}x$ for $z > 0$.

Lemma 3.1. *Let $z > 0, a > -1, b > -1, a + b = \alpha > -1$. Then*

$$\int_0^\infty \frac{x^a \mathrm{e}^{-x} \widetilde{p}_n(x)}{z+x} \mathrm{d}x = \mathrm{e}^z G^{2,1}_{2,3}\left(\begin{matrix} -n-b, a+n+1 \\ 0, a, -b \end{matrix} \middle| z \right).$$

Proof. By Theorem 3.1

$$\int_0^\infty \frac{x^a \mathrm{e}^{-x} \widetilde{p}_n(x)}{z+x} \mathrm{d}x = \int_\gamma \frac{\mathrm{d}u}{2\pi i} \frac{\Gamma(\alpha+n-u+1)\Gamma(u)}{\Gamma(n+1+u)\Gamma(\alpha+1-u)\Gamma(a+1-u)} \int_0^\infty \frac{x^{a-u} \mathrm{e}^{-x}}{z+x} \mathrm{d}x,$$

where the contour γ is chosen for this computation in such a way that $\Re u < 1+a$ in addition to $\Re u < 1+\alpha$ required by Theorem 3.1. Furthermore, we move the contour γ sufficiently to the right to ensure that $u = a$ is inside the contour. The change of the order of integration is justified because both single variable integrals are uniformly convergent on compact sets of their respective complementary variables and the original iterated integral is absolutely convergent. The inner integral is up to a normalization the Kummer function of the second kind (see [1] p. 194). A quick way of computing this integral is to substitute $\frac{1}{z+x} = \int_0^\infty \mathrm{e}^{-(z+x)s} \mathrm{d}s$ then switch the order of iterated integrals to get:

$$\int_0^\infty \frac{x^{a-u} \mathrm{e}^{-x}}{z+x} \mathrm{d}x = \mathrm{e}^z \Gamma(a+1-u) z^{a-u} \Gamma(u-a; z),$$

where $\Gamma(u-a;z)$ is the incomplete gamma function. To conclude this part of the computation one only needs to use $\Gamma(u-a;z) = \Gamma(u-a) + \gamma(u-a;z) = \Gamma(u-a) - \frac{z^{u-a}}{a-u}\ {}_1F_1(u-a, u-a+1; -z)$ which expresses the complementary incomplete gamma function γ in terms of the hypergeometric function ${}_1F_1$ (see [5] p. 266, formula (22)) thus obtaining

$$\int_0^\infty \frac{x^{a-u}\mathrm{e}^{-x}}{z+x}\mathrm{d}x = \mathrm{e}^z\Gamma(a+1-u)\Gamma(u-a)z^{a-u} - \mathrm{e}^z\Gamma(a-u)\ {}_1F_1(u-a, u-a+1; -z).$$

Putting the first term on the right side back into the contour integral gives the claim if one uses the shift formula eq. (A.2). To finish the proof we need to show that the contour integral of the second term vanishes. To this end we write explicitly the resulting contour integral obtaining, after omitting e^z and performing elementary computations with the gamma functions,

$$\int_\gamma \frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(\alpha+n-u+1)\Gamma(u)}{\Gamma(n+1+u)\Gamma(\alpha+1-u)}\sum_{j=0}\frac{1}{(u-a+j)j!}(-z)^j.$$

The series converges uniformly on γ hence to prove that this integral is zero it suffices to prove

$$\int_\gamma \frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(\alpha+n-u+1)\Gamma(u)}{\Gamma(n+1+u)\Gamma(\alpha+1-u)}\frac{1}{(u-a+j)} = 0, \quad j = 0, 1, \dots$$

The integrand is a meromorphic function with poles at $u = \{0, -1, \cdots, -n\} \cup \{a-j\}$ which, according to our choice of the contour γ, are all inside the contour. We can now extend this contour by adding a large circle with the center at $u = 0$ and radius r and observe that the integrand on such a large circle is, in view Lemma 2.2, $\mathcal{O}(u^{-2})$. This implies that in the limit of $r \to \infty$ the integral $\int_\gamma \frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(\alpha+n-u+1)\Gamma(u)}{\Gamma(n+1+u)\Gamma(\alpha+1-u)}\frac{1}{(u-a+j)} = [$ sum of residues of the integrand outside of $\gamma] = 0$.

□

The formula for $p_n^{(1)}(z)$ follows now easily from Lemma 3.1 and Theorem 3.2.

Theorem 3.3. *For $z \in \mathbb{C} \setminus \mathbb{R}_+$ the auxiliary functions of the first kind $p_n^{(1)}$ have the following Meijer G-function representation*

$$p_n^{(1)}(z) := \int_0^\infty \frac{p_n(x)x^a\mathrm{e}^{-x}\mathrm{d}x}{z-x} = \mathrm{e}^{-z}(-1)^{n+1}\pi_n G^{2,1}_{2,3}\left(\begin{matrix} -n-b, a+n+1 \\ 0, a, -b \end{matrix}\ \middle|\ -z\right).$$

A similar expression holds for $q_n^{(1)}$ by interchanging $a \leftrightarrow b$ and $\pi_n \leftrightarrow \eta_n$.

To see the relation between $p_n^{(1)}$ and p_n we formulate the following equivalent representation of $p_n^{(1)}$ whose gamma part is identical to that for p_n; see eq. (3-1)

Corollary 3.3. *For $z \in \mathbb{C} \setminus \mathbb{R}_+$ the auxiliary functions of the first kind admit an equivalent integral representation:*

$$p_n^{(1)}(z) = \mathrm{e}^{-z}\ (-1)^{n+1}\pi_n \int_\gamma \frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(1+n+\alpha-u)\Gamma(u)}{\Gamma(1+\alpha-u)\Gamma(n+1+u)\Gamma(1+a-u)}(-z)^{a-u}\frac{\pi}{\sin(\pi(u-a))} \tag{3-3}$$

Proof. The G-function occurring in the representation of p_n is $G^{1,1}_{2,3}\left(\begin{smallmatrix}-n+\alpha,n+1\\0,-\alpha,-a\end{smallmatrix}\middle| z\right)=G^{1,1}_{2,3}\left(\begin{smallmatrix}-n+\alpha,n+1\\0,-a,-\alpha\end{smallmatrix}\middle| z\right)$. Likewise, in view of Theorem 3.3 and the shift formula eq. (A.2) the G-function in the representation of $p_n^{(1)}$ can be written as $(-z)^a G^{2,1}_{2,3}\left(\begin{smallmatrix}-n-\alpha,n+1\\-a,0,-\alpha\end{smallmatrix}\middle| -z\right)$. Switching the first and the second coefficient in the bottom line of the latter G symbol can be done if one uses *Euler's reflection formula* to write $\Gamma(u-a)=\frac{\pi}{\Gamma(1+a-u)\sin(\pi(u-a))}$ which implies the claim. ☐

Remark 3.1. *It is instructive to verify that* $p_n^{(1)}(z)_+ - p_n^{(1)}(z)_- = -2\pi i\mathrm{e}^{-z}z^a p_n(z)$. *Indeed, let* $z>0$ *then*

$$(-z)_+^{a-u}-(-z)_-^{a-u}=z^{a-u}\left[e^{i\pi(u-a)}-e^{-i\pi(u-a)}\right]=z^{a-u}2i\sin(\pi(u-a)),$$

since the cut for z^x *is by convention (see A) along the negative real axis of* z. *Thus by the corollary above we get:*

$$\begin{aligned}&p_n^{(1)}(z)_+ - p_n^{(1)}(z)_- =\\&\mathrm{e}^{-z}\ (-1)^{n+1}\pi_n\int_\gamma\frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(n+\alpha+1-u)\Gamma(u)}{\Gamma(\alpha+1-u)\Gamma(n+1+u)\Gamma(a+1-u)}z^{a-u}\,2i\cancel{\sin(\pi(u-a))}\frac{\pi}{\cancel{\sin(\pi(u-a))}},\end{aligned}\tag{3-4}$$

which by Theorem 3.2 implies $p_n^{(1)}(z)_+ - p_n^{(1)}(z)_- = -2\pi i z^a e^{-z}p_n(z)$.

Theorem 3.4. *For* $z\in\mathbb{C}\setminus\mathbb{R}_-$ *the auxiliary functions of the second kind admit the* G*-function representation*

$$p_n^{(2)}(z) := -\int_0^\infty\int_0^\infty\frac{p_n(x)x^a y^b\mathrm{e}^{-x-y}\mathrm{d}x\mathrm{d}y}{(x+y)(z+y)} = (-1)^{n+1}\pi_n G^{3,1}_{2,3}\left(\begin{matrix}-n,\alpha+n+1\\0,\alpha,b\end{matrix}\middle| z\right)\tag{3-5}$$

Proof. Let us express $p_n^{(2)}$ in terms of $p_n^{(1)}$ as

$$p_n^{(2)}(z)=\int_0^\infty\frac{y^b e^{-y}}{z+y}p_n^{(1)}(-y)\mathrm{d}y.$$

Since $p_n^{(1)}(-y)=(-1)^{n+1}\pi_n\,\mathrm{e}^y\,G^{2,1}_{2,3}\left(\begin{smallmatrix}-n-b,a+n+1\\a,0,-b\end{smallmatrix}\middle| y\right)$ we obtain

$$p_n^{(2)}(z)=(-1)^{n+1}\pi_n\int_0^\infty\frac{y^b}{z+y}G^{2,1}_{2,3}\left(\begin{matrix}-n-b,a+n+1\\a,0,-b\end{matrix}\middle| y\right)\mathrm{d}y=(-1)^{n+1}\pi_n\,G^{3,1}_{2,3}\left(\begin{matrix}-n,\alpha+n+1\\0,\alpha,b\end{matrix}\middle| z\right),$$

by eq. (A.4) and eq. (A.2). ☐

Again to see the relation between $p_n^{(1)}$ and $p_n^{(2)}$ we establish an equivalent representation

Corollary 3.4. *For* $z\in\mathbb{C}\setminus\mathbb{R}_-$ *the auxiliary functions of the second kind admit an equivalent integral representation:*

$$p_n^{(2)}(z)=(-1)^{n+1}\pi_n\int_\gamma\frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(u)\Gamma(u+a)\Gamma(1+n+b-u)}{\Gamma(1+b-u)\Gamma(a+n+1+u)}z^{b-u}\frac{\pi}{\sin(\pi(u-b))}.$$

Proof. Using the shift formula given by eq. (A.2) we write $G^{3,1}_{2,3}\left({-n,\alpha+n+1 \atop 0,\alpha,b}\,\middle|\,z\right) = z^b G^{3,1}_{2,3}\left({-n-b,a+n+1 \atop -b,a,0}\,\middle|\,z\right)$. On the other hand, in view of Theorem 3.3, the G function in the representation of $p_n^{(1)}$ is $G^{2,1}_{2,3}\left({-n-b,a+n+1 \atop 0,a,-b}\,\middle|\,-z\right)$. Switching the first and the last coefficient in the bottom line of the G symbol can be done if one uses *Euler's reflection formula* to write $\Gamma(u-b) = \frac{\pi}{\Gamma(1+b-u)\sin(\pi(b-u))}$ which implies the claim. □

Remark 3.2. *By analogy with the* $p_n^{(1)}$ *case we easily verify the jump of* $p_n^{(2)}(z)$ *for* $z<0$ *to be* $p_n^{(2)}(z)_+ - p_n^{(2)}(z)_- = 2\pi i(-z)^b \mathrm{e}^z p_n^{(1)}(z)$. *Indeed, let* $z<0$ *then*

$$(z)_+^{b-u} - (z)_-^{b-u} = (-z)^{b-u}\left[e^{i\pi(b-u)} - e^{-i\pi(b-u)}\right] = (-z)^{b-u} 2i\sin(\pi(b-u)),$$

since the cut for z^x *is by convention (see A) along the negative real axis of* z*. Thus, again, by the corollary above we get:*

$$p_n^{(2)}(z)_+ - p_n^{(2)}(z)_- = (-z)^b(-1)^{n+1}\pi_n G^{2,1}_{2,3}\left(\begin{matrix}-n-b, a+n+1\\ 0, a, -b\end{matrix}\,\middle|\,-z\right),$$

which by Theorem 3.3 implies $p_n^{(2)}(z)_+ - p_n^{(2)}(z)_- = 2\pi i(-z)^b \mathrm{e}^z p_n^{(1)}(z)$.

3.2 The kernels for finite n

In [3] we proved a variety of generalizations of Christoffel–Darboux identities. In this section, however, we show that in the case at hand it is possible to compute the associated kernels directly and in an elementary way from the formulæ above. We will begin with $K_{00}^{(n)}$ as defined in eq. (2-10).

Theorem 3.5. *The principal kernel* $K_{00}^{(n)}(x,y)$ *is given by*

$$K_{00}^{(n)}(x,y) = \int_0^1 t^\alpha G_{a,n}(tx) G_{b,n}(ty), \qquad \textit{where} \tag{3-6}$$

$$G_{c,n}(x) := \int_\gamma \frac{\mathrm{d}u}{2i\pi} \frac{x^{-u}}{\Gamma(c+1-u)} \frac{\Gamma(u)\Gamma(n+\alpha+1-u)}{\Gamma(n+u)\Gamma(\alpha-u+1)} = G^{1,1}_{2,3}\left(\begin{matrix}-\alpha-n, n\\ 0, -c, -\alpha\end{matrix}\,\middle|\,x\right). \tag{3-7}$$

Proof. We start with writing the formula for the kernel in terms of G-functions. By Theorem 3.2 $K_{00}^{(n)}(x,y) = \sum_{j=0}^{n-1}(2j+\alpha+1) G^{1,1}_{2,3}\left({-\alpha-j,j+1 \atop 0,-\alpha,-a}\,\middle|\,x\right) G^{1,1}_{2,3}\left({-\alpha-j,j+1 \atop 0,-\alpha,-b}\,\middle|\,y\right)$, while Christoffel-Darboux kernel for Jacobi polynomials is given by $K_n(x,y) = \sum_{j=0}^{n-1}(2j+\alpha+1) G^{1,1}_{2,2}\left({-\alpha-j,j+1 \atop 0,-\alpha}\,\middle|\,x\right) G^{1,1}_{2,2}\left({-\alpha-j,j+1 \atop 0,-\alpha}\,\middle|\,y\right)$ (see Propositions 2.2 and 2.3). This means that the integral representations obtained for $K_n(x,y)$ are only modified by extra factors $\frac{1}{\Gamma(1+a-u)}, \frac{1}{\Gamma(1+b-v)}$ coming from the additional third index appearing in the bottom of the respective G-symbols of $\widetilde{p}_j(x), \widetilde{q}_n(y)$ which proves the claim. □

Remark 3.3. *We can write explicitly the double integral representation of the kernel by copying the analogous formula from Theorem 2.2 and attaching additional factors* $\frac{1}{\Gamma(1+a-u)}, \frac{1}{\Gamma(1+b-v)}$ *as explained in the proof above. The formula reads:*

$$K_{00}^{(n)}(x,y) = \\ \int_{\gamma^2} \frac{\mathrm{d}u}{2\pi i}\frac{\mathrm{d}v}{2\pi i}\frac{x^{-u}y^{-v}}{\Gamma(a+1-u)\Gamma(b+1-v)(1+\alpha-v-u)}\frac{\Gamma(u)\Gamma(v)\Gamma(n+\alpha+1-u)\Gamma(n+\alpha+1-v)}{\Gamma(n+u)\Gamma(n+v)\Gamma(\alpha-u+1)\Gamma(\alpha-v+1)}. \tag{3-8}$$

Note that this integral representation, as opposed to the one appearing in 2.2, is valid for any $x, y \neq 0$. *Observe also that the function* $G_{c,n}(x) = G^{1,1}_{2,3}\left(\begin{smallmatrix}-\alpha-n,n\\0,-c,-\alpha\end{smallmatrix}\middle| x\right)$ *is a natural generalization of the function* $\widehat{G}_n(x) = G^{1,1}_{2,2}\left(\begin{smallmatrix}-\alpha-n,n\\0,-\alpha\end{smallmatrix}\middle| x\right)$ *known from Theorem 2.2.*

We are now ready to compute the remaining kernels. Using Lemma 2.1 we find

Theorem 3.6. *The kernel* $K_{10}^{(n)}(x,y)$ *is given by*

$$K_{10}^{(n)}(x,y) = x^a \mathrm{e}^x \int_0^1 t^\alpha \widetilde{G}_{a,n}(tx)G_{b,n}(ty), \quad \textit{where} \tag{3-9}$$

$$\widetilde{G}_{c,n}(x) := \int_\gamma \frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(u-c)\Gamma(u)\Gamma\left(n+\alpha+1-u\right)}{\Gamma(\alpha+1-u)\Gamma\left(n+u\right)}x^{-u} = G^{2,1}_{2,3}\left(\begin{matrix}-\alpha-n,n\\0,-c,-\alpha\end{matrix}\middle| x\right). \tag{3-10}$$

Likewise, by symmetry, the kernel $K_{01}^{(n)}(x,y)$ *is given by:*

$$K_{01}^{(n)}(x,y) = y^b \mathrm{e}^y \int_0^1 t^\alpha G_{a,n}(tx)\widetilde{G}_{b,n}(ty) \tag{3-11}$$

$$\tag{3-12}$$

with $\widetilde{G}_{c,n}$ *defined in eq. (3-10) and* $G_{c,n}$ *defined in eq. (3-7).*

Proof. From the definition of $K_{10}^{(n)}$ in eq. (2-12), Theorem 3.3 and eq. (A.2) we obtain

$$K_{10}^{(n)}(x,y) := -\sum_{j=0}^{n-1} p_j^{(1)}(-x)q_j(y) = \\ x^a\mathrm{e}^x\sum_{j=0}^{n-1}(2j+\alpha+1)G^{2,1}_{2,3}\left(\begin{matrix}-\alpha-j,j+1\\0,-a,-\alpha\end{matrix}\middle| x\right)G^{1,1}_{2,3}\left(\begin{matrix}-\alpha-j,j+1\\0,-\alpha,-b\end{matrix}\middle| y\right).$$

We observe that the dependence on j is identical to the one covered in Lemma 2.1. Hence by that lemma

$$K_{10}^{(n)}(x,y) = x^a\mathrm{e}^x\left\{\int_0^1 t^\alpha G^{2,1}_{2,3}\left(\begin{matrix}-\alpha-n,n\\0,-a,-\alpha\end{matrix}\middle| tx\right)G^{1,1}_{2,3}\left(\begin{matrix}-\alpha-n,n\\0,-\alpha,-b\end{matrix}\middle| ty\right)\mathrm{d}t- \\ \int_0^1 t^\alpha G^{1,0}_{0,1}\left(\begin{matrix}\\-a\end{matrix}\middle| tx\right)G^{0,0}_{0,1}\left(\begin{matrix}\\-b\end{matrix}\middle| ty\right)\mathrm{d}t\right\} = x^a\mathrm{e}^x\int_0^1 t^\alpha G^{2,1}_{2,3}\left(\begin{matrix}-\alpha-n,n\\0,-a,-\alpha\end{matrix}\middle| tx\right)G^{1,1}_{2,3}\left(\begin{matrix}-\alpha-n,n\\0,-\alpha,-b\end{matrix}\middle| ty\right)\mathrm{d}t,$$

since the second term, or more precisely $G^{0,0}_{0,1}\left(\begin{smallmatrix}\\-b\end{smallmatrix}\middle| ty\right)$, vanishes by eq. (A.5). □

Likewise,

Theorem 3.7. *The Kernel* $K_{11}^{(n)}(x,y)$ *is given by*

$$K_{11}^{(n)}(x,y) = x^a y^b \mathrm{e}^{x+y} \int_0^1 \widetilde{G}_{a,n}(tx)\widetilde{G}_{b,n}(ty) t^{\alpha} \mathrm{d}t - \frac{\mathrm{e}^{x+y}}{x+y} \tag{3-13}$$

with $\widetilde{G}_{c,n}$ *defined in eq. (3-10).*

Proof. We plug the expressions for $p_n^{(1)}, q_n^{(1)}$ of Theorem 3.3 into the definition of $K_{11}^{(n)}$ (2-14) to get

$$K_{11}^{(n)}(x,y) + \frac{1}{x+y} := \sum_{j=0}^{n-1} p_j^{(1)}(-x) q_j^{(1)}(-y) =$$
$$x^a y^b \mathrm{e}^{x+y} \sum_{j=0}^{n-1} (2j+\alpha+1) G^{2,1}_{2,3}\left(\begin{matrix} -\alpha-j, j+1 \\ 0, -a, -\alpha \end{matrix} \middle| x \right) G^{2,1}_{2,3}\left(\begin{matrix} -\alpha-j, j+1 \\ 0, -b, -\alpha \end{matrix} \middle| y \right).$$

Again, the j-dependence of this expression is covered by Lemma 2.1 and applying it we obtain:

$$K_{11}^{(n)}(x,y) + \frac{1}{x+y} = x^a y^b \mathrm{e}^{x+y} \{ \int_0^1 t^{\alpha} G^{2,1}_{2,3}\left(\begin{matrix} -\alpha-n, n \\ 0, -a, -\alpha \end{matrix} \middle| tx \right) G^{2,1}_{2,3}\left(\begin{matrix} -\alpha-n, n \\ 0, -b, -\alpha \end{matrix} \middle| ty \right) \mathrm{d}t -$$
$$\int_0^1 t^{\alpha} G^{1,0}_{0,1}\left(\begin{matrix} \\ -a \end{matrix} \middle| tx \right) G^{1,0}_{0,1}\left(\begin{matrix} \\ -b \end{matrix} \middle| ty \right) \mathrm{d}t \}.$$

By inspection $G^{1,0}_{0,1}\left(\begin{matrix} \\ -a \end{matrix} \middle| tx \right) = (tx)^{-a} \mathrm{e}^{-tx}$, hence an elementary computation gives the final answer

$$K_{11}^{(n)}(x,y) + \frac{1}{x+y} = x^a y^b \mathrm{e}^{x+y} \{ \int_0^1 t^{\alpha} G^{2,1}_{2,3}\left(\begin{matrix} -\alpha-n, n \\ 0, -a, -\alpha \end{matrix} \middle| tx \right) G^{2,1}_{2,3}\left(\begin{matrix} -\alpha-n, n \\ 0, -b, -\alpha \end{matrix} \middle| ty \right) \mathrm{d}t +$$
$$\frac{1}{x^a y^b} \frac{(\mathrm{e}^{-(x+y)} - 1)}{x+y} \},$$

which implies the claim. □

Remark 3.4. *Similar to the integral formula for the kernel* K_{00}^n *(see eq. (3-8)) all the remaining kernels have analogous integral representations in terms of the double path integral, all involving the kernel* $\frac{1}{1+\alpha-u-v}$.

3.3 The Meijer-G random point field

In order to describe the statistics of the Cauchy matrix model near $x = 0 = y$ we recall the definition of the kernels for the correlation functions, that is, the K and H kernels defined by eq. (1-6), eq. (1-5)) respectively.

3.3.1 Asymptotics of the kernels $K_{\mu\nu}^{(n)}$

From the explicit integral expressions for the functions $G_{c,n}, \widetilde{G}_{c,n}$ and the kernels in Theorems 3.5, 3.6, 3.7, we shall now derive the behaviour under the rescaling

$$x = \frac{\zeta}{n^2}\ , \qquad y = \frac{\xi}{n^2}, \quad n \to \infty. \tag{3-14}$$

Remark 3.5. *It should be mentioned here that if we had started with a model in the form*

$$\mathrm{d}\mu(M_1, M_2) = \frac{1}{\mathcal{Z}_n}\mathrm{d}M_1\mathrm{d}M_2\frac{(\det M_1)^a(\det M_2)^b\mathrm{e}^{-n\mathrm{Tr}(M_1+M_2)}}{\det(M_1+M_2)^n} \tag{3-15}$$

then the relevant rescaling would have been $x \mapsto xn^{-3}$, because the n scaling in the exponential is simply absorbed by a rescaling $\widetilde{x} = nx$.

Theorem 3.8 (Kernels in the asymptotic regime)**.** *The following asymptotic estimates for $n \to \infty$ hold uniformly over compact subsets of the independent variables*

$$n^{-\alpha-1}G_{c,n}(\zeta n^{-2}) = \overbrace{\int_\gamma \frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(u)}{\Gamma(c+1-u)\Gamma(\alpha+1-u)}\zeta^{-u}}^{=:G_c(\zeta)} + \mathcal{O}(n^{-1}), \tag{3-16}$$

$$n^{-\alpha-1}\widetilde{G}_{c,n}(\zeta n^{-2}) = \underbrace{\int_\gamma \frac{\mathrm{d}u}{2\pi i}\frac{\Gamma(u-c)\Gamma(u)}{\Gamma(\alpha+1-u)}\zeta^{-u}}_{=:\widetilde{G}_c(\zeta)} + \mathcal{O}(n^{-1}), \tag{3-17}$$

or in terms of G-functions

$$n^{-\alpha-1}G^{1,1}_{2,3}\left(\begin{matrix}-\alpha-n, n\\ 0, -c, -\alpha\end{matrix}\,\middle|\,\zeta n^{-2}\right) = G^{1,0}_{0,3}\left(\begin{matrix}\\ 0, -c, -\alpha\end{matrix}\,\middle|\,\zeta\right) + \mathcal{O}(n^{-1}),$$

$$n^{-\alpha-1}G^{2,1}_{2,3}\left(\begin{matrix}-\alpha-n, n\\ 0, -c, -\alpha\end{matrix}\,\middle|\,\zeta n^{-2}\right) = G^{2,0}_{0,3}\left(\begin{matrix}\\ 0, -c, -\alpha\end{matrix}\,\middle|\,\zeta\right) + \mathcal{O}(n^{-1}).$$

Correspondingly, the asymptotic behavior of the four kernels is

$$n^{-2\alpha-2}K_{00}^{(n)}(\zeta n^{-2}, \xi n^{-2}) \to \int_0^1 G_a(t\zeta)G_b(t\xi)t^\alpha\mathrm{d}t, \tag{3-18}$$

$$n^{-2a-2}K_{01}^{(n)}(\zeta n^{-2}, \xi n^{-2}) \to \xi^b\int_0^1 G_a(t\zeta)\widetilde{G}_b(t\xi)t^\alpha\mathrm{d}t, \tag{3-19}$$

$$n^{-2b-2}K_{10}^{(n)}(\zeta n^{-2}, \xi n^{-2}) \to \zeta^a\int_0^1 \widetilde{G}_a(t\zeta)G_b(t\xi)t^\alpha\mathrm{d}t, \tag{3-20}$$

$$n^{-2}K_{11}^{(n)}(\zeta n^{-2}, \xi n^{-2}) \to \zeta^a\xi^b\int_0^1 \widetilde{G}_a(t\zeta)\widetilde{G}_b(t\xi)t^\alpha\mathrm{d}t - \frac{1}{\zeta+\xi}, \tag{3-21}$$

and the convergence of the above limits is all within $\mathcal{O}(n^{-1})$.

Proof. The proof of this theorem is immediate from the Stirling approximation formula in Lemma 2.2, the explicit integral expressions for $G_{c,n}, \widetilde{G}_{c,n}$ in eq. (3-7), eq. (3-10) and the convolution form for the kernels in Theorems 3.5, 3.6, 3.7. $\square$

Now we turn our attention to the H kernels defined in eq. (1-5). Using the rescaling (3-14) and the asymptotics of the kernels in Theorem 3.8 we find the following expressions

$$\lim_{n\to\infty} n^{-2} H_{00}^{(n)}(x,y) = \zeta^a \xi^b \int_0^1 G_a(t\zeta) G_b(t\xi) t^\alpha \mathrm{d}t, \tag{3-22}$$

$$\lim_{n\to\infty} n^{-2} H_{01}^{(n)}(x,y) = \zeta^a \xi^b \int_0^1 G_a(t\zeta) \widetilde{G}_b(t\xi) t^\alpha \mathrm{d}t, \tag{3-23}$$

$$\lim_{n\to\infty} n^{-2} H_{10}^{(n)}(x,y) = \zeta^a \xi^b \int_0^1 \widetilde{G}_a(t\zeta) G_b(t\xi) t^\alpha \mathrm{d}t, \tag{3-24}$$

$$\lim_{n\to\infty} n^{-2} H_{11}^{(n)}(x,y) = \zeta^a \xi^b \int_0^1 \widetilde{G}_a(t\zeta) \widetilde{G}_b(t\xi) t^\alpha \mathrm{d}t - \frac{1}{\zeta+\xi}. \tag{3-25}$$

Remark 3.6. *The scaling factor n^{-2} in front of the kernels $H_{\mu\nu}^{(n)}$ is natural since kernels transform as $\sqrt{\mathrm{d}x\mathrm{d}y}$.*

The limiting kernels above define a novel class of random point fields.

Proposition 3.1 (Meijer-G random point field and universal class)**.** *In the scaling limit the correlations of the eigenvalues of M_1, M_2 are determined by the two–level random point field with kernels*

$$\mathcal{G}_{00}(\zeta,\xi) = \zeta^a \xi^b \int_0^1 G_a(t\zeta) G_b(t\xi) t^\alpha \mathrm{d}t, \qquad \mathcal{G}_{01}(\zeta,\xi) = \zeta^a \xi^b \int_0^1 G_a(t\zeta) \widetilde{G}_b(t\xi) t^\alpha \mathrm{d}t, \tag{3-26a}$$

$$\mathcal{G}_{10}(\zeta,\xi) = \zeta^a \xi^b \int_0^1 \widetilde{G}_a(t\zeta) G_b(t\xi) t^\alpha \mathrm{d}t, \qquad \mathcal{G}_{11}(\zeta,\xi) = \zeta^a \xi^b \int_0^1 \widetilde{G}_a(t\zeta) \widetilde{G}_b(t\xi) t^\alpha \mathrm{d}t - \frac{1}{\zeta+\xi}, \tag{3-26b}$$

where $\xi, \zeta \in \mathbb{R}_+$.

Remark 3.7. *It is a nontrivial fact that the correlation functions defined using the kernels in Proposition 3.1 are all positive and define an actual random point field: of course this is hard to see directly from the definition of the kernels; it nevertheless follows from the fact that this is the limit of a well–defined point field.*

If we absorb the powers in the definition of $H_a, \widetilde{H}_a$ as in Definition 1.1 then the formulas becomes even more symmetric.

Corollary 3.5. *In the scaling limit the correlations of the eigenvalues of M_1, M_2 are determined by the two–level random point field with kernels*

$$\mathcal{G}_{00}(\zeta,\xi) = \int_0^1 H_a(t\zeta) H_b(t\xi) \mathrm{d}t, \qquad \mathcal{G}_{01}(\zeta,\xi) = \int_0^1 H_a(t\zeta) \widetilde{H}_b(t\xi) \mathrm{d}t, \tag{3-27a}$$

$$\mathcal{G}_{10}(\zeta,\xi) = \int_0^1 \widetilde{H}_a(t\zeta) H_b(t\xi) \mathrm{d}t, \qquad \mathcal{G}_{11}(\zeta,\xi) = \int_0^1 \widetilde{H}_a(t\zeta) \widetilde{H}_b(t\xi) \mathrm{d}t - \frac{1}{\zeta+\xi}, \tag{3-27b}$$

where $\xi, \zeta \in \mathbb{R}_+$ and the kernels are given by the Meijer G functions

$$H_c(z) = G^{1,0}_{0,3}\left(\begin{matrix} \\ c, 0, -\alpha + c \end{matrix} \middle| z \right), \quad \widetilde{H}_c(z) = G^{2,0}_{0,3}\left(\begin{matrix} \\ c, 0, -\alpha + c \end{matrix} \middle| z \right).$$

Thus we have completed the proof of Theorem 1.2 stated in the introduction.

Remark 3.8. *It is easy to see by direct computation that the functions $H_c, \tilde{H}_c$ are connected to hypergeometric functions ${}_0F_2$. For example*

$$H_a(z) = \frac{z^a \; {}_0F_2\left({}_{1+a, 1+\alpha}; -z\right)}{\Gamma\left(1+a\right)\Gamma\left(1+\alpha\right)}, \tag{3-28}$$

while $\widetilde{H}_a(z)$ is, for $a \neq 0, 1, \cdots$, a linear combination of two hypergeometric functions of type ${}_0F_2$

$$\widetilde{H}_a(z) = \frac{\Gamma\left(a\right) \; {}_0F_2\left({}_{1+b, 1-a}; z\right)}{\Gamma\left(1+b\right)} + \frac{\Gamma\left(-a\right) z^a \; {}_0F_2\left({}_{1+a, 1+\alpha}; z\right)}{\Gamma\left(1+\alpha\right)}. \tag{3-29}$$

The remaining cases $a = n = 0, 1, \cdots$ are handled by taking limits $a \to n$. For example

$$\widetilde{H}_0(z) = -\frac{\ln z}{\Gamma(1+b)} + \mathcal{O}(1) \qquad \text{if } a = 0.$$

3.3.2 Expressions of the kernels in "integrable" form

We note that the functions $H_a(t), \widetilde{H}_a(t)$ solve differential equations of the third order; this fact is implied by their identification as Meijer-G functions. A simple derivation is as follows: denote by $A(u)$ the rational expression of Γ functions in the integrand of H_c or $\widetilde{H}_c$; then a direct inspection shows

$$A(u+1) = \mp u(u+c)(\alpha - c - u)A(u) \quad \Rightarrow \quad \Delta_\zeta(\Delta_\zeta + \alpha - c)(\Delta_\zeta - c)f(\zeta) = \mp\zeta f(\zeta), \tag{3-30}$$

where $\Delta_\zeta = \zeta\frac{d}{d\zeta}$, the upper sign is for H_c, the lower for $\widetilde{H}_c$ respectively. Now, let $c \in \{a, b\}$ and denote by $f_t(\zeta) := f(t\zeta)$, $g_t(\xi) := g(t\xi)$, and let f be one of H_a or $\widetilde{H}_a$ and g one of H_b or $\widetilde{H}_b$. Since Δ is scale invariant we have

$$g_t(\xi)\Delta_\zeta(\Delta_\zeta + b)(\Delta_\zeta - a)f_t(\zeta) = \mp t\zeta f_t(\zeta)g_t(\xi), \tag{3-31a}$$

$$f_t(\zeta)\Delta_\xi(\Delta_\xi + a)(\Delta_\xi - b)g_t(\xi) = \mp t\xi f_t(\zeta)g_t(\xi). \tag{3-31b}$$

Adding these two equations together, dividing by t and using that $\Delta_\zeta f_t(\zeta) = \Delta_t f_t(\zeta)$ (and similarly for $g_t(\xi)$) we find

$$\partial_t\left[g_t\Delta_t^2 f_t - \Delta_t g_t \Delta_t f_t + f_t\Delta_t^2 g_t + (b-a)(g_t\Delta_t f - f_t\Delta_t g_t) - ab f_t g_t\right] = \mp\zeta + \mp\xi)f_t g_t. \tag{3-32}$$

Integrating with respect to t from 0 to $0 < \tau$ we obtain

$$\mp \zeta + \mp \xi) \int_{0+}^{\tau} f_t g_t \mathrm{d}t = f_t \Delta_t^2 g_t - \Delta_t f_t \Delta_t g_t + g_t \Delta_t^2 f_t + (b-a)\left(g_t \Delta_t f_t - f_t \Delta_t g\right) - ab\, f_t g_t \Big|_{0+}^{\tau} = \\ \left[f(t\zeta), \Delta_\zeta f(t\zeta), \Delta_\zeta^2 f(t\zeta)\right] \begin{bmatrix} -ab & a-b & 1 \\ b-a & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} g(t\xi) \\ \Delta_\xi g(t\xi) \\ \Delta_\xi^2 g(t\xi) \end{bmatrix}_{t=0+}^{\tau} , \tag{3-33}$$

where in the last line we used again the scale invariance of Δ. The last computation motivates the following definition

Definition 3.1. *Let $f = f(\zeta)$ and $g = g(\xi)$. Then the* **point-split bilinear concomitant** *is defined as*

$$\mathcal{B}(f,g) := f\Delta_\xi^2 g - (\Delta_\zeta f)(\Delta_\xi g) + g\Delta_\zeta^2 f + (b-a)\left(g\Delta_\zeta f - f\Delta_\xi g\right) - ab\, fg = \\ \left[f, \Delta_\zeta f, \Delta_\zeta^2 f\right] \begin{bmatrix} -ab & a-b & 1 \\ b-a & -1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} g \\ \Delta_\xi g \\ \Delta_\xi^2 g \end{bmatrix} . \tag{3-34}$$

If $\zeta = \pm\xi$, suitably chosen to make the left side of eq. (3-33) vanish, then eq. (3-33) implies $\partial_\xi \mathcal{B}(f,g) = 0$. This is a form of the *bilinear concomitant* [7] for equations in duality, which explains our naming convention.

From this point onward we will be interested in $\tau = 1$. The evaluation of the right hand side of eq. (3-33) at $t = 0+$ is case dependent.

Case H_a, H_b. If $f = H_a,\ g = H_b$ then the evaluation at $t = 0$ vanishes: indeed by eq. (3-28)

$$H_c(\zeta) = \zeta^c \left(\frac{1}{\Gamma(1+c)\Gamma(1+\alpha)} + \mathcal{O}(\zeta) \right) \tag{3-35}$$

and it suffices therefore to verify $\mathcal{B}(\zeta^a, \xi^b) = 0$ and

$$\lim_{t \to 0+} \mathcal{B}((t\zeta)^{a+k}, (t\xi)^{b+l}) = \lim_{t \to 0+} t^{\alpha+k+l} \mathcal{B}(\zeta^{a+k}, \xi^{b+l}) = 0, \text{ for } 1 \le k+l \ ,$$

since $\alpha + 1 > 0$.

Case $\widetilde{H}_a, H_b$ or vice versa. This case is similar to the previous case; one uses eq. (3-29) and analyzes cases. The relevant points to remember are that by our assumptions $a+1 > 0, b+1 > 0$ and $\alpha + 1 > 0$. In all cases $\mathcal{B}$ vanishes at $t = 0+$.

Case $\widetilde{H}_a, \widetilde{H}_b$. In this case the $\lim_{t\to 0+} \mathcal{B}(\widetilde{H}_a, \widetilde{H}_b)(t\zeta, t\xi) = -1$. This follows from the fact that in the expansion of $\widetilde{H}_a$ (or $\widetilde{H}_b$) there is one more pairing that the previous cases did not have. This is the pairing of the leading term in eq. (3-29), namely in $\frac{\Gamma(a)\ {}_0F_2\left({}_{1+b,1-a};z\right)}{\Gamma(1+b)}$, with its analog in $\widetilde{H}_b$. Extracting the leading term leads to $\mathcal{B}(\frac{\Gamma(a)}{\Gamma(1+b)}, \frac{\Gamma(b)}{\Gamma(1+a)}) = -ab\frac{\Gamma(a)\Gamma(b)}{\Gamma(1+a)\Gamma(1+b)} = -1$.

It is now elementary to substitute the above results into eq. (3-33) and, further, into the definition of $\mathcal{G}$ kernels given by eq. (3-27a).

Proposition 3.2. *Let $\mathcal{B}$ be the point-split bilinear concomitant defined by 3.1. Then the correlation kernels of the Meijer-G random point field satisfy*

$$\mathcal{G}_{00}(\zeta,\xi) = -\frac{\mathcal{B}(H_a(\zeta), H_b(\xi))}{\zeta+\xi}, \qquad \mathcal{G}_{01}(\zeta,\xi) = \frac{\mathcal{B}(H_a(\zeta), \widetilde{H}_b(\xi))}{-\zeta+\xi}, \tag{3-36a}$$

$$\mathcal{G}_{10}(\zeta,\xi) = \frac{\mathcal{B}(\widetilde{H}_a(\zeta), H_b(\xi))}{\zeta-\xi}, \qquad \mathcal{G}_{11}(\zeta,\xi) = \frac{\mathcal{B}(\widetilde{H}_a(\zeta), \widetilde{H}_b(\xi))}{\zeta+\xi}. \tag{3-36b}$$

A The Meijer-G functions

For the convenience of the reader we recall the definition of the Meijer-G function

$$G^{m,n}_{p,q}\left(\begin{matrix} a_1,\ldots,a_p \\ b_1,\ldots,b_q\end{matrix}\ \middle|\ z\right) = \int_\gamma \frac{\mathrm{d}u}{2\pi i}\, \frac{\prod_{j=1}^m \Gamma(b_j+u)\prod_{j=1}^n \Gamma(1-a_j-u)}{\prod_{j=m+1}^q \Gamma(1-b_j-u)\prod_{j=n+1}^p \Gamma(a_j+u)}\, z^{-u}. \tag{A.1}$$

The contour γ is a contour that leaves the poles of the $\Gamma(b_j+u)$'s to the left and the poles of the $\Gamma(1-a_j-u)$'s to the right (the implicit assumption is that none of the poles of the former coincides with any of the poles of the latter). The contour extends to infinity in with $|\arg(u)| < \frac{\pi}{2}$ (in the right half plane) if $p > q$, while it extends to infinity with $\frac{\pi}{2} < \arg(u) < \frac{3\pi}{2}$ (in the left half plane) if $p \le q$. It is the latter case that is relevant for us. We assume therefore that the branch of z^{-u} is chosen in such a way that the cut is for $z < 0$. There is an extensive literature on G-functions (see e.g. [8, 9]). We collect several useful formulas used in the paper.

1. Shifting formula:
$$z^\sigma G^{m,n}_{p,q}\left(\begin{matrix}\mathbf{a_p}\\ \mathbf{b_q}\end{matrix}\ |\, z\right) = G^{m,n}_{p,q}\left(\begin{matrix}\mathbf{a_p}+\sigma\\ \mathbf{b_q}+\sigma\end{matrix}\ \middle|\ z\right), \tag{A.2}$$
where $+\sigma$ means shifting every entry by σ.

2. Integrals containing products of two G-functions:
$$\int_0^\infty G^{m,n}_{p,q}\left(\begin{matrix}\mathbf{a_p}\\ \mathbf{b_q}\end{matrix}\ \middle|\ \frac{x}{\eta}\right) G^{\mu,\nu}_{\sigma,\tau}\left(\begin{matrix}\mathbf{c}_\sigma\\ \mathbf{d}_\tau\end{matrix}\ \middle|\ \omega x\right) dx = \eta\, G^{n+\mu,\, m+\nu}_{q+\sigma,\, p+\tau}\left(\begin{matrix}-b_1,\ldots,-b_m,\mathbf{c}_\sigma,-b_{m+1},\ldots,-b_q\\ -a_1,\ldots,-a_n,\mathbf{d}_\tau,-a_{n+1},\ldots,-a_p\end{matrix}\ \middle|\ \eta\omega\right). \tag{A.3}$$
Since $G^{1,1}_{1,0}\left(\begin{matrix}a\\ b\end{matrix}\ \middle|\ -z\right)$ is proportional to $z^b\ {}_1F_0(1+b-a,-z) = z^b\frac{1}{(1+z)^{1+b-a}}$ we obtain [8]
$$\int_0^\infty \frac{y^{\alpha-1}}{(z+y)^\sigma} G^{m,n}_{p,q}\left(\begin{matrix}\mathbf{a_p}\\ \mathbf{b_q}\end{matrix}\ \middle|\ \mu y\right) \mathrm{d}y = \frac{z^{\alpha-\sigma}}{\Gamma(\sigma)} G^{m+1,n+1}_{p+1,q+1}\left(\begin{matrix}1-\alpha,\mathbf{a_p}\\ \sigma-\alpha,\mathbf{b_q}\end{matrix}\ \middle|\ \mu z\right). \tag{A.4}$$

3. For $p \leq q$

$$G^{0,n}_{p,q}\left(\begin{matrix}\mathbf{a_p}\\ \mathbf{b_q}\end{matrix} \mid z\right) = 0. \tag{A.5}$$

B Correlation functions

In this appendix we recall the definitions and formulas used to study the correlation functions for generic Cauchy matrix models (no restrictions on the measures). We use the notation that is slightly different than the one used in [2] to accommodate the needs of the present paper. Given two measures $d\alpha(x) = \alpha(x)\mathrm{d}x$, $\mathrm{d}\beta(y) = \beta(y)$ with densities $\alpha(x), \beta(y)$ respectively, both supported on the positive half-line $\mathbb{R}_+$, the Cauchy matrix model is the probability measure on the pairs (M_1, M_2) of positive semi-definite $N \times N$ matrices given by

$$\mathrm{d}\mu(M_1, M_2) = \frac{\alpha(M_1)\beta(M_2)\mathrm{d}M_1\mathrm{d}M_2}{\mathcal{Z}_N \det(M_1+M_2)^N}.$$

where $\alpha(M)$ (or $\beta(B)$) stands for the induced measure on the spectrum of M, i.e. $\prod_j \alpha(x_j)$ where x_j are the (positive) eigenvalues of M. Let us introduce two families of polynomials $\{p_j(x), j = 0, 1, ..\}$ and $\{q_j(y), j = 0, 1, ..\}$ which are biorthonormal with respect to the pairing with the Cauchy kernel

$$\iint_{\mathbb{R}_+^2} \frac{p_j(x)\alpha(x)q_k(y)\beta(y)}{x+y}\mathrm{d}x\mathrm{d}y = \delta_{j,k},$$

with the technical proviso that the leading coefficients are identical and positive to render all polynomials unique. Following [2], but changing slightly the notation, we introduce four kernels used in the present paper

$$K_{00}^{(n)}(x,y) := \sum_{j=0}^{n-1} p_j(x)q_j(y), \qquad K_{01}^{(n)}(x,y) := \int K_{00}^{(n)}(x',y)\frac{\alpha(x')\mathrm{d}x'}{x+x'},$$
$$K_{10}^{(n)}(x,y) := \int K_{00}^{(n)}(x,y')\frac{\beta(y')\mathrm{d}y'}{y+y'}, \quad K_{11}^{(n)}(x,y) := \iint K_{00}^{(n)}(x',y')\frac{\alpha(x')\mathrm{d}x'\beta(y')\mathrm{d}y'}{(x+x')(y+y')} - \frac{1}{x+y}.$$

With this notation in place the correlation functions for r eigenvalues $x_1, \ldots, x_r$ of M_1 and s eigenvalues $y_1, \ldots, y_s$ of M_2 can be shown to be given by

$$\mathcal{R}^{(r,s)}(x_1,\ldots,x_r;y_1,\ldots,y_s) = \prod_{j=1}^r \alpha(x_j)\,\mathrm{d}x_j \prod_{k=1}^s \beta(y_k)\mathrm{d}y_k \det\left[\begin{array}{c|c} \left[K_{01}^{(n)}(x_i,x_j)\right]_{i,j\leq r} & \left[K_{00}^{(n)}(x_i,y_j)\right]_{i\leq r, j\leq s} \\ \hline \left[K_{11}^{(n)}(y_i,x_j)\right]_{i\leq s, j\leq r} & \left[K_{10}^{(n)}(y_i,y_j)\right]_{i,j\leq s}\end{array}\right] \tag{B.1}$$

It is easy to write the correlation function as one determinant by including the products in front of the determinant above as determinants of diagonal matrices. This is not a unique procedure but the choice that works naturally for the present paper is the following. We write symbolically

$$\mathcal{R}^{(r,s)}(x_1,\ldots,x_r;y_1,\ldots,y_s) = \det\left(\begin{bmatrix}\alpha(\mathbf{x}) & 0\\ 0 & \mathbf{I}\end{bmatrix}\begin{bmatrix}\mathbf{K_{01}}(\mathbf{x},\mathbf{x}) & \mathbf{K_{00}}(\mathbf{x},\mathbf{y})\\ \mathbf{K_{11}}(\mathbf{y},\mathbf{x}) & \mathbf{K_{10}}(\mathbf{y},\mathbf{y})\end{bmatrix}\begin{bmatrix}\mathbf{I} & 0\\ 0 & \beta(\mathbf{y})\end{bmatrix}\right),$$

where $\alpha(\mathbf{x}) = \mathrm{diag}(\alpha(x_1), \cdots, \alpha(x_r)), \mathrm{diag}(\beta(y_1), \cdots, \beta(y_s))$ respectively. Hence

$$\mathcal{R}^{(r,s)}(x_1, \ldots, x_r; y_1, \ldots, y_s) = \det \left(\begin{bmatrix} \alpha(\mathbf{x})\mathbf{K_{01}}(\mathbf{x}, \mathbf{x}) & \alpha(\mathbf{x})\mathbf{K_{00}}(\mathbf{x}, \mathbf{y})\beta(\mathbf{y}) \\ \mathbf{K_{11}}(\mathbf{y}, \mathbf{x}) & \mathbf{K_{10}}(\mathbf{y}, \mathbf{y})\beta(\mathbf{y}) \end{bmatrix} \right).$$

This leads to the definition of new kernels:

$$H_{00}^{(n)}(x,y) := \alpha(x)\beta(y)K_{00}^{(n)}(x,y), \qquad H_{01}^{(n)}(x,y) := \alpha(x)K_{01}^{(n)}(x,y), \tag{B.2a}$$

$$H_{10}^{(n)}(x,y) := \beta(y)K_{10}^{(n)}(x,y), \qquad H_{11}^{(n)}(x,y) := K_{11}^{(n)}(x,y). \tag{B.2b}$$

References

[1] R. Beals and R. Wong. *Special functions*, volume 126 of *Cambridge Studies in Advanced Mathematics.* Cambridge University Press, Cambridge, 2010. A graduate text.

[2] M. Bertola, M. Gekhtman, and J. Szmigielski. The Cauchy two–matrix model. *Comm. Math. Phys.*, 287(3):983–1014, 2009.

[3] M. Bertola, M. Gekhtman, and J. Szmigielski. Cauchy biorthogonal polynomials. *J. Approx. Theory*, 162(4):832–867, 2010.

[4] A. Borodin. Biorthogonal ensembles. *Nuclear Phys. B*, 536(3):704–732, 1999.

[5] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Higher transcendental functions. Vol. I.* Robert E. Krieger Publishing Co. Inc., Melbourne, Fla., 1981.

[6] J. L. Fields. The asymptotic expansion of the Meijer G-function. *Math. Comp.*, 26:757–765, 1972.

[7] E. L. Ince. *Ordinary Differential Equations.* Dover Publications, New York, 1944.

[8] Y. L. Luke. *The special functions and their approximations, Vol. I.* Mathematics in Science and Engineering, Vol. 53. Academic Press, New York, 1969.

[9] Y. L. Luke. *The special functions and their approximations. Vol. II.* Mathematics in Science and Engineering, Vol. 53. Academic Press, New York, 1969.

[10] G. Szegö. *Orthogonal polynomials.* American Mathematical Society Colloquium Publications, Vol. 23. Revised ed. American Mathematical Society, Providence, R.I., 1959.

[11] C. A. Tracy and H. Widom. Level spacing distributions and the Bessel kernel. *Comm. Math. Phys.*, 161(2):289–309, 1994.

[12] C. A. Tracy and H. Widom. Level-spacing distributions and the Airy kernel. *Comm. Math. Phys.*, 159(1):151–174, 1994.